



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

March 26, 2015

Alfred Gwilliam
Denka Registrations BV
4 Tar Rock Rd
Westport CT 06880

Subject: Label Amendment – Methomyl Fly Bait Mitigation Language
Product Name: Lurectron Scatterbait
EPA Registration Number: 7319-6
Application Date: March 11, 2015
Decision Number: 502738

Dear Mr. Gwilliam:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 7319-6
Decision No. 502738

with FIFRA section 6. If you have any questions, please contact Carlyn Petrella by phone at 703-347-0439, or via email at petrella.carlyn@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Gebken", with a long horizontal flourish extending to the right.

Richard Gebken
Product Manager 10
Invertebrate & Vertebrate Branch 2
Office of Pesticide Programs

LURECTRON® SCATTERBAIT

With Muscalure Fly Attractant

Insecticide for use against flies in commercial poultry houses, livestock barns, feed lots, canneries, beverage producers, meat, poultry and food processing plants, and on walkways in caged layer houses.


ACTIVE INGREDIENTS

Methomyl [CAS No. 16752-77-5]	1.00%
Z-9-Tricosene [CAS No. 27519-02-4]	0.26%
INERT INGREDIENTS	98.74%
TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION

KEEP OUT OF REACH OF CHILDREN
See Back Panel for Additional Cautions

EPA Registration No. 7319-6
Denka International b.v.
Hanzeweg 1
3771 NG Barneveld
The Netherlands



EPA Establishment No. 7319-NL-1
NON-REFILLABLE CONTAINER

Net Contents ___lb ___ Gm



Do not use any form of methomyl fly bait product around or inside homes, or any other place where children are likely to be present.
It is illegal to use this product with the intention to kill raccoons, skunks, opossums, coyotes, wolves, dogs, cats, or any other non-target species.

FIRST AID

If swallowed:

- ▶ Call a poison control center or doctor immediately for treatment advice.
- ▶ Have the person sip a glass of water if able to swallow.
- ▶ Do not induce vomiting unless told to do so by a poison control center or doctor.
- ▶ Do not give anything by mouth to an unconscious person.

If on skin or clothing:

- ▶ Take off contaminated clothing.
- ▶ Rinse skin immediately with plenty of water for 15 - 20 minutes.
- ▶ Call a poison control center or doctor immediately for treatment advice.

If in eyes:

- ▶ Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.
- ▶ Remove contact lenses, if present, after first 5 minutes, then continue rinsing the eye.
- ▶ Call a poison control center or doctor immediately for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Centre at Tel: 1-800-858-7378 for emergency medical treatment information.
NOTE TO PHYSICIAN: Methomyl is a cholinesterase inhibitor; atropine is antidotal.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin or clothing.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long sleeved shirt and long pants.
- Socks and shoes.
- Chemical resistant gloves made of waterproof material.

Follow manufacturer's instructions for cleaning / maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, the user should wash thoroughly and change into clean clothing.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to birds, fish, aquatic vertebrates and mammals.

Collect, cover or incorporate granules spilled on soil surface. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighbouring areas. Do not contaminate water when disposing of equipment wash-water or rinsate. Some of the chemical in this product are known to leach through soil into groundwater under certain conditions of label use. Use of this chemical in areas where the soil is permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS OF USE

It is a violation of federal law to use this product in a manner inconsistent with its labelling. Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Use this product only for the sites, pests, and application methods described on this label.

Do not apply around or inside homes or any other place where children are likely to be present. Use only in areas inaccessible to children, pets, and food producing animals. Insecticide Resistance Management Farnam DIE FLY™ can be an important tool in the overall management of housefly populations. Incorporate an Integrated Pest Management program which includes: sanitation, proper manure management, elimination of breeding areas, physical barriers, liquid spray treatments, fly traps (mechanical, light, glue), and periodic rotation of baits with different modes of action.

For use as scatter baits: Use around the outside of the following commercial establishments where children are unlikely to be present: feed lots, poultry houses, livestock barns, canneries, beverage plants, meat and poultry processing establishments, foods processing plants and commercial refuse dumpsters which are fenced or otherwise enclosed. Do not use around commercial dumpsters that are not enclosed. Apply only to dumpsters that have a lid or can be closed. Dumpster doors and lids must be closed after applying bait. Do not use around dumpsters in residential settings. May be used on walkways in caged layer houses. Do not treat areas where animals can feed on bait. Do not apply directly to water. Keep children, pets, and livestock clear from scatter bait treatment area. Distribute bait from container or use an implement to avoid handling the bait directly. Scatter bait evenly at a rate of ¼ lb (4 oz.) product per 500 square feet of area. Do not apply in piles. Replenish every 3 to 5 days or when bait is no longer visible in the treated area.

For use in bait stations: Use in any commercial bait station secured at least 4 feet above the ground and in areas not accessible to children, pets, and livestock. Use around the outside of feed lots, poultry houses, livestock barns, canneries, beverage plants, meat and poultry processing establishments, foods processing plants, kennels, and refuse dumpsters associated with these establishments. May be used in fenced or otherwise enclosed areas inaccessible to children and pets around the outside of restaurants, supermarkets, and bakeries. May be used inside caged layer houses. Pour 1 oz.(28 g) into each bait station and use at the rate of 2 bait stations per 500 square feet of fly feeding area. Replenish every 3 to 5 days or when bait is no longer visible in the bait station. When refilling it may be necessary to first dispose of dead flies and any product remaining in bait station as described below under Storage and Disposal. Replace bait station if physically damaged.

For use as a paste: Mix ¼ lb (4 oz.) product with ½ cup (4 oz.) water stirring well to form a paste. Apply paste with a brush on walls, windowsills, and support beams where flies can land on the outside of livestock barns and poultry houses such that children, pets, and livestock cannot come into contact with paste. Reapply every 3 to 5 days of when flies no longer appear to be affected by landing on treated area.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage and disposal. Open dumping is prohibited.

Pesticide storage: Store in cool, dry, secure place and keep container tightly closed. Store container away from children and pets.

Pesticide disposal: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

Container handling:

Non refillable Container. DO NOT reuse or refill this container. After completely emptying container into application equipment, offer for recycling, if available, or reconditioning, if appropriate, or puncture

and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Bait station disposal

The bait station is reusable. When the container becomes cracked or broken, offer for recycling or dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

WARRANTY STATEMENT

Manufacturer makes no warranty, express or implied, concerning the use of this product other than as indicated on the label. Buyer assumes all risks of use, storage and handling of this material when such use and handling are contrary to label directions.